



FEEDLOT CP BALANCER RU1200 MEDICATED GRANULAR

For improved feed efficiency for cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Monensin 1,200 g/ton

GUARANTEED ANALYSIS

Calcium (Ca)	28.0% Min. ... 32.0% Max.
Salt (NaCl)	13.0% Min. ... 15.0% Max.
Potassium (K)	3.0% Min.
Copper (Cu)	750 ppm Min.
Selenium (Se)	8.8 ppm Min.
Zinc (Zn)	3,000 ppm Min.
Vitamin A	120,000 IU/lb. Min.
Vitamin D-3	12,000 IU/lb. Min.
Vitamin E	120 IU/lb. Min.

INGREDIENTS

Calcium Carbonate, Salt, Potassium Chloride, Mineral Oil, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Thiamine Mononitrate.

FEEDING DIRECTIONS

Feed continuously to cattle being fed in confinement for slaughter at the rate of **.25** to **.33** pound per head per day, to provide 150 to 200 mg of Monensin daily. Consult your **PEET'S** Representative for ration balancing services and medicated options available.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

NOTICE: Expiration date of 30 days after manufacture.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.