



## FEEDLOT NATURAL RU400 MEDICATED PELLET

For improved feed efficiency in cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

### ACTIVE DRUG INGREDIENTS

Monensin ..... 400 g/ton

### GUARANTEED ANALYSIS

Crude Protein	32.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	10.0% Max.
Calcium (Ca)	7.0% Min. ... 8.0% Max.
Phosphorus (P)	0.85% Min.
Salt (NaCl)	2.5% Min. ... 3.5% Max.
Potassium (K)	1.5% Min.
Selenium (Se)	2.2 ppm Min.
Vitamin A	40,000 IU/lb. Min.
Vitamin D-3	4,000 IU/lb. Min.
Vitamin E	40 IU/lb. Min.

### INGREDIENTS

Soybean Meal, Distillers Dried Grains/Solubles, Calcium Carbonate, Pork Meat & Bone Meal, Cane Molasses, Salt, Linseed Meal, Dehydrated Alfalfa Meal, Wheat Middlings, Pellet Coating, Potassium Chloride, Dried Aspergillus oryzae Fermentation Product, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Thiamine Mononitrate.

### FEEDING DIRECTIONS

Feed 1 pound per head per day to feedlot cattle to provide 200 mg of Monensin daily.

Consult your **PEET'S** Representative for ration balancing services to provide the optimum usage of grain and roughage.

---

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

---

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feed are safe for use in cattle and goats only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

