



CALF STARTER BT-60 MEDICATED

For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* for cattle weighing up to 800 pounds and for increased rate of gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers) when fed continuously.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Lasalocid 60 g/ton

GUARANTEED ANALYSIS

Crude Protein	18.0% Min.
Crude Fat	4.0% Min.
Crude Fiber	7.0% Max.
Acid Detergent Fiber	9.0% Max.
Calcium (Ca)	0.7% Min. ... 1.2% Max.
Phosphorus (P)	0.5% Min.
Salt (NaCl)	.3% Min. ... 0.8% Max.
Potassium (K)	0.8% Min.
Selenium (Se)	0.4 ppm. Min.
Vitamin A	15,000 IU/lb. Min.
Vitamin D-3	3,000 IU/lb. Min.
Vitamin E	30 IU/lb. Min.

INGREDIENTS

Rolled Yellow Corn, Whole Oats, Cane Molasses, Soybean Meal, Linseed Meal, Pork Meat and Bone Meal, Distillers Dried Grains/Solubles, Calcium Carbonate, Choice White Grease, Salt, Lignin Sulfonate, Dicalcium Phosphate, Dried Aspergillus oryzae Fermentation Product, Sodium Selenite, MOL Enhancer, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Natural and Artificial Flavors.

FEEDING DIRECTIONS

Feed continuously at the rate of 1.5 pounds per 100 lbs. of body-weight daily for control of coccidiosis in calves up to 800 pounds. For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers) feed at the rate of 2 to 10 pounds daily to provide 60 to 300 mg Lasalocid daily.

Calf Weight	LBS. to Feed Daily	LASALOCID Provided (mg)
100	1.5	45
200	3.0	90
300	4.5	135
400	6.0	180
500	7.5	225
600	9.0	270
700	10.5	315
800	12.0	360

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal. **CAUTION:** The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration that could be fatal to cattle and sheep.

