



SPRINT 10-15 DC MEDICATED PELLET

Designed to be fed as the sole ration to early weanling pigs until they reach 15 pounds of body weight. Recommended as a creep feed when sow milk production is inadequate or sows are nursing large litters.

NET WEIGHT 50 LBS. (22.7kg)

For treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline and for control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin when fed as directed on tag.

ACTIVE DRUG INGREDIENTS

Chlortetracycline 400 grams/ton
 Tiamulin Hydrogen Fumerate 35 grams/ton

GUARANTEED ANALYSIS

Crude Protein	22.0% Min.
Lysine	155% Min.
Crude Fat	8.0% Min.
Crude Fiber	2.0% Max
Calcium (Ca)	0.8% Min. ... 1.0% Max.
Phosphorus (P)	.8% Min.
Selenium (Se)	.3 ppm Min.
Zinc (Zn)	2500 ppm Min.
Vitamin A	6,800 IU/lb. Min.

INGREDIENTS

Ground Yellow Corn, Dried Whey, Rolled Oat Groats, Soybean Meal, Porcine Animal Plasma, Soybean Oil, Fish Meal, Lactose, Dicalcium Phosphate, Soy Protein Concentrate, Calcium Carbonate, Choice White Grease, Zinc Oxide, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Artificial Flavor, L-Lysine, Sodium Selenite, DL-Methionine, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D-3), Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Mineral Oil, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydriodide, Dried Aspergillus oryzae Fermentation Product, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Sodium Saccharin, Manganese Amino Acid Complex.

See WARNING on Label

Consult your PEET'S Representative for specific feeding recommendations.

WARNING: Withdraw 2 days prior to slaughter.

CAUTION: Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

SPECIAL CONSIDERATIONS: Protect medicated feed from excessive temperatures. Do not store at warm temperatures 104° F (40° C), for more than 7 days.

LIMITATIONS FOR USE: Feed continuously as sole ration for 14 days. Not for use in swine weighing more than 250 pounds. Feed approximately 400 g/ton Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day.

