



ULTRA SPRINT 5-10 DC MEDICATED PELLET

Designed to be fed as sole ration to early weaning pigs until they reach 10 lbs. of body weight.

NET WEIGHT 50 LBS. (22.7kg)

For treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline and for control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin when fed as directed on tag.

ACTIVE DRUG INGREDIENTS

Chlortetracycline 400 grams/ton

Tiamulin hydrogen fumarate 35 grams/ton

GUARANTEED ANALYSIS

Crude Protein	23.0% Min.
Lysine	1.8% Min.
Crude Fat	10.0% Min.
Crude Fiber	2.0% Max
Calcium (Ca)	0.85% Min. ... 1.05% Max.
Phosphorus (P)	.85% Min.
Selenium (Se)	.3 ppm Min.
Zinc (Zn)	2500 ppm Min.
Vitamin A	6,500 IU/lb. Min.

INGREDIENTS

Rolled Oat Groats, Dried Whey, Lactose, Soybean Oil, Porcine Animal Plasma, Soybean Meal, Fish Meal, Ground Yellow Corn, Soy Protein Concentrate, Dicalcium Phosphate, Calcium Carbonate, Zinc Oxide, L-Lysine, Animal Fat with BHA Preservative, Lecithin, Casein, DL-Methionine, Artificial Flavor, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Sodium Selenite, Dried *Aspergillus oryzae* Fermentation Product, Sodium Saccharin, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D-3), Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Mineral Oil, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydriodide, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Zinc Amino Acid Complex, Manganese Amino Acid Complex.

DIRECTIONS FOR USE

Consult your **PEET'S** Representative for specific feeding recommendations.

WARNING: Withdraw 2 days prior to slaughter.

CAUTION: Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds. **SPECIAL CONSIDERATIONS:** Protect medicated feed from excessive temperatures. Do not store at warm temperatures, 104° F (40° C), for more than 7 days. **LIMITATIONS FOR USE:** Not for use in swine weighing more than 250 pounds. Feed approximately 400 g/ton Chlortetracycline varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight per day.

