



## CALF SUPPLEMENT BT 200 MEDICATED PELLET

Controls coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* when mixed and fed to cattle.

**NET WEIGHT 50 LBS. (22.7kg)**

### ACTIVE DRUG INGREDIENT

Lasalocid ..... 200 g/ton

### GUARANTEED ANALYSIS

Crude Protein	38.0% Min.
Crude Fat	3.4% Min.
Crude Fiber	8.0% Max.
Acid Detergent Fiber	12.0% Max.
Calcium (Ca)	1.75% Min. ... 2.25% Max.
Phosphorus (P)	1.1% Min.
Salt (NaCl)	0.75% Min. ... 1.25% Max.
Potassium (K)	1.5% Min.
Selenium (Se)	1.4 ppm Min.
Vitamin A	50,000 IU/lb. Min.
Vitamin D-3	10,000 IU/lb. Min.
Vitamin E	100 IU/lb. Min.

### INGREDIENTS

Soybean Meal, Cottonseed Meal, Linseed Meal, Pork Meat and Bone Meal, Distillers Dried Grains/Solubles, Cane Molasses, Calcium Carbonate, Choice White Grease, Salt, Dicalcium Phosphate, Dried Aspergillus oryzae Fermentation Product, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide, Natural and Artificial Flavors.

### FEEDING DIRECTIONS

Mix into the daily ration to provide 0.45 pounds of supplement per hundred pounds of bodyweight making the Lasalocid intake 45 mg per hundred pounds of bodyweight daily.

	18% RATION	16% RATION
Corn	1000 lbs.	1250 lbs.
Oats	400 lbs.	300 lbs.
<b>CALF SUPP. BT 200</b>	600 lbs.	450 lbs.
<b>TOTAL</b>	<b>2000 lbs.</b>	<b>2000 lbs.</b>
Feeding Rate		
PER 100# BWT	1.5 lbs.	2.0 lbs.

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Consult your **PEET'S** Representative for ration balancing services available. **USE LIMITATIONS:** Feed only to cattle weighing up to 800 pounds providing a maximum of 360 mg of Lasalocid per head per day. Not for free choice feeding. **CAUTION:** The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration that could be fatal to cattle and sheep.

