



FEEDLOT 32/16 RU500 MEDICATED PELLET

For improved feed efficiency for cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENTS

Monensin 500 g/ton

GUARANTEED ANALYSIS

Crude Protein (not more than 16.0% equivalent crude protein from non-protein nitrogen.)	32.0% Min.
Crude Fat	2.2% Min.
Crude Fiber	10.5% Max.
Calcium (Ca)	10.0% Min. .. 12.0% Max.
Phosphorus (P)	0.5% Min.
Salt (NaCl)	2.5% Min. .. 3.5% Max.
Potassium (K)	1.0% Min.
Selenium (Se)	1.1 ppm Min.
Vitamin A	40,000 IU/lb. Min.
Vitamin D-3	4,000 IU/lb. Min.
Vitamin E	40 IU/lb. Min.

INGREDIENTS

Wheat Middlings, Calcium Carbonate, Linseed Meal, Soybean Meal, Pork Meat & Bone Meal, Urea, Dehydrated Alfalfa Meal, Cane Molasses, Distillers Dried Grains/Solubles, Salt, Sodium Bentonite, Pellet Coating, Potassium Chloride, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Zinc Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Thiamine Mononitrate, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

FEEDING DIRECTIONS

Feed continuously at the rate of 1 pound per head per day to beef cattle being fed in confinement to provide 250 mg of Monensin and 50 mg of thiamin daily. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Expiration date of 30 days after manufacture.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.
