



## TRIPLE 12 OXY2 MINERAL MEDICATED

NET WEIGHT 50 LBS. (22.7kg)

### ACTIVE DRUG INGREDIENT

Oxytetracycline ..... 2.0 grams/lb.

### GUARANTEED ANALYSIS

Calcium (Ca)	12.0% Min. ... 14.0% Max
Phosphorus (P)	12.0% Min.
Salt (NaCl)	12.0% Min. ... 14.0% Max.
Magnesium (Mg)	2.0% Min.
Potassium (K)	1.0% Min.
Copper (Cu)	1,200 ppm Min.
Selenium (Se)	25 ppm Min.
Zinc (Zn)	4,000 ppm Min.
Vitamin A	200,000 IU/lb. Min.
Vitamin D-3	20,000 IU/lb. Min.
Vitamin E	200 IU/lb. Min.

### INGREDIENTS

Dicalcium Phosphate, Salt, Monocalcium Phosphate, Distillers Dried Grains/Solubles, Potassium/Magnesium Sulfate, Magnesium Oxide, Sodium Selenite, Cane Molasses, Mineral Oil, Potassium Chloride, Sodium Sesquicarbonate, Yeast Culture, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Ethylenediamine Dihydrochloride, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Natural and Artificial Flavors, Artificial Color.

### FEEDING DIRECTIONS

For the prevention and treatment of the early stages of shipping fever complex at the rate of 4 ounces per head daily for 3-5 days before and after arrival to growing beef cattle to provide 0.5 grams of Terramycin daily. Consult your **PEET'S** Representative for ration balancing services to provide the optimum usage of grain and roughage.

**WARNING:** Withdraw 5 days before slaughter.