



CO-PRODUCT BREEDER MINERAL R1600 MEDICATED GRANULAR

A complex complete macro mineral, trace mineral and vitamin supplement for cattle designed to balance rations containing corn processing by-products.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Monensin 1,600 g/ton

GUARANTEED ANALYSIS

Calcium (Ca)	19.0% Min. ... 22.0% Max.
Phosphorus (P)	2.0% Min.
Salt (NaCl)	12.0% Min. ... 14.0% Max.
Magnesium (Mg)	2.0% Min.
Potassium (K)	1.0% Min.
Copper (Cu)	3,000 ppm Min.
Selenium (Se)	25 ppm Min.
Zinc (Zn)	4,000 ppm Min.
Vitamin A	300,000 IU/lb. Min.
Vitamin D-3	30,000 IU/lb. Min.
Vitamin E	300 IU/lb. Min.
Thiamine	800 mg/lb. Min.

INGREDIENTS

Calcium Carbonate, Salt, Distillers Dried Grains/Solubles, Monocalcium Phosphate, Dicalcium Phosphate, Potassium/Magnesium Sulfate, Magnesium Oxide, Sodium Selenite, Cane Molasses, Mineral Oil, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Ethylenediamine Dihydriodide, Potassium Chloride, Sodium Sesquicarbonate, Yeast Culture, Thiamine, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Natural and Artificial Flavors, Artificial Color.

MIXING & FEEDING DIRECTIONS

Mix 3 ounces in not less than 1 pound of feed and either hand feed or mix into the total ration to provide 150 mg Monensin per head per day. During the first 5 days, cattle should receive no more than 100 mg per head per day. Contact your **PEET'S** Representative for specific rations and information.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. This product should not be offered to sheep due to the copper level. Feed as the sole source of selenium. Do not free choice.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.
