



## CATTLE SUPREME 8 CTC2.6 MINERAL MEDICATED GRANULAR

Controls active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline.

NET WEIGHT 50 LBS. (22.7kg)

### ACTIVE DRUG INGREDIENT

Chlortetracycline ..... 2.6 grams/lb.

### GUARANTEED ANALYSIS

Calcium (Ca)	14.0% Min. ... 16.0% Max.
Phosphorus (P)	8.0% Min.
Salt (NaCl)	18.0% Min. ... 20.0% Max.
Magnesium (Mg)	2.0% Min.
Potassium (K)	0.5% Min.
Copper (Cu)	2,200 ppm Min.
Selenium (Se)	26.4 ppm Min.
Zinc (Zn)	6,000 ppm Min.
Vitamin A	300,000 IU/lb. Min.
Vitamin D-3	30,000 IU/lb. Min.
Vitamin E	300 IU/lb. Min.

### INGREDIENTS

Monocalcium Phosphate, Salt, Calcium Carbonate, Dicalcium Phosphate, Distillers Dried Grains/Solubles, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Cane Molasses, Soybean Oil, Potassium/Magnesium Sulfate, Potassium Chloride, Yeast Culture, Ethylenediamine Dihydriodide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Natural and Artificial Flavors, Artificial Color.

### FEEDING DIRECTIONS

Feed at the rate listed below to beef cattle under 700 pounds to provide 350 mg of Chlortetracycline daily and .5 mg/lb. of bodyweight per day to beef cattle over 700 pounds. Consult your PEET'S Representative for ration balancing services.

Cattle Bodyweight (lbs.)	Ounces/Head/Day
600-700	2.2
800	2.5
1,000	3.1
1,200	3.7
1,400	4.3

**WARNING:** Withdrawal 48 hours prior to slaughter. Do not use in calves to be processed for veal.