



FEEDLOT PERFORMER RU1200 MEDICATED PELLET

For improved feed efficiency for cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Monensin 1,200 g/ton

GUARANTEED ANALYSIS

Crude Protein	10.0% Min.
Crude Fat	2.4% Min.
Crude Fiber	12.0% Max.
Calcium (Ca)	10.0% Min. ... 12.0% Max.
Phosphorus	2.0% Min.
Salt (NaCl)	4.5% Min. ... 5.5% Max.
Potassium (K)	2.5% Min.
Selenium (Se)	6.6 ppm Min.
Vitamin A	160,000 IU/lb. Min.
Vitamin D-3	16,000 IU/lb. Min.
Vitamin E	160 IU/lb. Min.

INGREDIENTS

Wheat Middlings, Calcium Carbonate, Distillers Dried Grains/Solubles, Soybean Hulls, Dicalcium Phosphate, Salt, Potassium Chloride, Cane Molasses, Choice White Grease, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

FEEDING DIRECTIONS

Feed at the rate of 1/3 - 1/2 pound per head per day to provide 200 to 300 mg of Monensin daily. Feed continuously to growing feedlot cattle being fed in confinement for slaughter to supplement high protein hay or haylage. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.
