



MGA PAC 1 MEDICATED PELLET

For increased rate of weight gain, improved feed efficiency and suppression of estrus (heat) in heifers fed for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENTS

Melengestrol Acetate 1.0 mg/lb.

GUARANTEED ANALYSIS

Crude Protein	7.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	18.0% Max.
Calcium (Ca)	14.0% Min. ... 16.0% Max.
Phosphorus (P)	0.2% Min.
Salt (NaCl)	4.5% Min. ... 5.4% Max.
Potassium (K)	0.5% Min.

INGREDIENTS

Calcium Carbonate, Wheat Middlings, Soybean Hull, Salt, Cane Molasses, Choice White Grease.

FEEDING DIRECTIONS

Feed to heifers at the rate of 1/2 pound per head per day to provide 0.5 mg Melengestrol Acetate daily.

CAUTION: Withdrawal periods of 3 to 5 days should be avoided to prevent the possibility that heifers may come into estrus (heat) at loading time. Not effective in steers and spayed heifers. Heifers intended for reproduction: Do not exceed 24 days of feeding. A reduced conception rate can be expected if heifers are bred at estruses observed within 1 to 12 days after withdrawal of Melengestrol Acetate, whereas heifers bred at subsequent observed estrus are expected to have normal conception rates. Consult your **PEET'S** Representative for ration balancing services to provide the optimum usage of grain and roughage.