



## MGA PAC 1 MEDICATED PELLET

For increased rate of weight gain, improved feed efficiency and suppression of estrus (heat) in heifers fed for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

### ACTIVE DRUG INGREDIENTS

Melengestrol Acetate ..... 1.0 mg/lb.

### GUARANTEED ANALYSIS

Crude Protein	7.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	18.0% Max.
Calcium (Ca)	14.0% Min. ... 16.0% Max.
Phosphorus (P)	0.2% Min.
Salt (NaCl)	4.5% Min. ... 5.4% Max.
Potassium (K)	0.5% Min.

### INGREDIENTS

Calcium Carbonate, Wheat Middlings, Soybean Hull, Salt, Cane Molasses, Choice White Grease.

### FEEDING DIRECTIONS

Feed to heifers at the rate of 1/2 pound per head per day to provide 0.5 mg Melengestrol Acetate daily.

**CAUTION:** Withdrawal periods of 3 to 5 days should be avoided to prevent the possibility that heifers may come into estrus (heat) at loading time. Not effective in steers and spayed heifers. Heifers intended for reproduction: Do not exceed 24 days of feeding. A reduced conception rate can be expected if heifers are bred at estruses observed within 1 to 12 days after withdrawal of Melengestrol Acetate, whereas heifers bred at subsequent observed estrus are expected to have normal conception rates. Consult your **PEET'S** Representative for ration balancing services to provide the optimum usage of grain and roughage.