



COXY CURB MEDICATED PELLET

An aid in the prevention or treatment of coccidiosis in beef and dairy calves caused by *Eimeria bovis* and *Eimeria zuernii*.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Amprovine 1.25%

INGREDIENTS

Wheat Middlings, Soybean Hulls, Calcium Carbonate, Dehydrated Alfalfa Meal, Cane Molasses, Choice White Grease.

FEEDING DIRECTIONS

21 DAY PREVENTION PROGRAM: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, mix 0.04 pound into the feed for each 100 pounds of calf weight daily to provide 227 mg Amprovine per 100 lbs. bodyweight per day for 21 days.

CALF WEIGHT	COXY CURB lbs./100 Head Daily	
	Prevention	Treatment
200 lbs.	8	16
300 lbs.	12	24
400 lbs.	16	32
500 lbs.	20	40
600 lbs.	24	48

5 DAY TREATMENT PROGRAM: When one or more calves show signs of coccidiosis, it is likely that the entire herd has been exposed. Treat on a herd basis by mixing 0.08 pound into the feed for each 100 pounds of calf weight daily to provide 454 mg Amprovine per 100 lbs. bodyweight per day for 5 days.

CAUTION: Do not use Amprovine in feeds containing bentonite.

WARNING: Withdraw 24 hours before slaughter. Use as sole source of Amprolium. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.
