



FEEDLOT NATURAL BT400 MEDICATED PELLET

For improved feed efficiency in cattle fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENTS

Lasalocid 400 grams/ton

GUARANTEED ANALYSIS

Crude Protein	32.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	10.0% Max.
Calcium (Ca)	7.0% Min. ... 8.0% Max.
Phosphorus (P)	0.85% Min.
Salt (NaCl)	2.5% Min. ... 3.5% Max.
Potassium (K)	1.5% Min.
Selenium (Se)	2.2 ppm Min.
Vitamin A	40,000 IU/lb. Min.
Vitamin D-3	4,000 IU/lb. Min.
Vitamin E	40 IU/lb. Min.

INGREDIENTS

Soybean Meal, Calcium Carbonate, Distillers Dried Grains/Solubles, Pork Meat & Bone Meal, Cane Molasses, Salt, Linseed Meal, Dehydrated Alfalfa Meal, Wheat Middlings, Pellet Coating, Dicalcium Phosphate, Dried Aspergillus oryzae Fermentation Product, Potassium Chloride, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

FEEDING DIRECTIONS

Feed 1 pound per head per day continuously to feedlot beef cattle to provide 200 mg of Lasalocid daily. Consult your **PEET'S** Representative for ration balancing services to provide the optimum usage of grain and roughage and other medicated options available.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

CAUTION: The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration, which could be fatal to cattle and sheep.

