



FEEDLOT 40/28 RU 500 MEDICATED PELLET

For improved feed efficiency for cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

2 ton minimum

ACTIVE DRUG INGREDIENTS

Monensin 500 g/ton

GUARANTEED ANALYSIS

Crude Protein (includes not more than 28% equivalent crude protein from non-protein nitrogen.)	40.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	10.5% Max.
Calcium (Ca)	10.0% Min. ... 12.0% Max.
Phosphorus (P)	0.5% Min.
Salt (NaCl)	2.5% Min. ... 3.5% Max.
Potassium (K)	1.0% Min.
Selenium (Se)	3.2 ppm Min.
Vitamin A	40,000 IU/lb. Min.
Vitamin D-3	4,000 IU/lb. Min.
Vitamin E	40 IU/lb. Min.

INGREDIENTS

Wheat Middlings, Calcium Carbonate, Distillers Dried Grains/Solubles, Urea, Dehydrated Alfalfa Meal, Cane Molasses, Salt, Soybean Meal, Pork Meat & Bone Meal, Pellet Coating, Magnesium Oxide, Potassium Chloride, Amino Resin, Calcium Sulfate, Potassium/Magnesium Sulfate, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

DIRECTIONS FOR USE

Feed 1 pound per head per day continuously to cattle being fed in confinement for slaughter to provide 250 mg of Monensin daily. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage and other medicated options available.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.
