



CTC 10 MEDICATED

For the treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Chlortetracycline 10 grams/lb.

GUARANTEED ANALYSIS

Crude Protein	10.0% Min.
Crude Fat	2.4% Min.
Crude Fiber	18.0% Max.
Calcium (Ca)	3.5% Min. ... 4.5% Max.
Phosphorus (P)	0.4% Min.
Potassium (K)	0.8% Min.

INGREDIENTS

Wheat Middlings, Soybean Hulls, Calcium Carbonate, Dehydrated Alfalfa Meal, Cane Molasses, Choice White Grease.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. Withdraw 48 hours prior to slaughter when feeding .5 mg or more per pound of bodyweight daily or when feeding 350 mg or more per head per day.

FEEDING DIRECTIONS

CALVES

For increased rate of weight gain and improved feed efficiency.

BODYWEIGHT	LEVEL OF CTC per head daily	FEEDING RATE Pounds per 100 Head
*Up to 250 lb.	0.1 mg/lb. of bwt	Up to .25 lb.
250 to 400 lb.	25 - 70 mg	.25-.7 lb.

*LIMITATIONS FOR USE: In milk replacers or starter feeds

GROWER CATTLE

For increased rate of weight gain, improved feed efficiency & reduction of liver condemnations due to liver abscesses.

BODYWEIGHT	LEVEL OF CTC per head daily	FEEDING RATE Pounds per 100 Head
Over 400 lb.	70 mg	.7 lb.

Beef Cattle

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline.

LEVEL OF CTC per head daily	FEEDING RATE Pounds per 100 Head
350 mg	3.5 lb.

10 MG/LB. OF BWT. DAILY

For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline when fed to calves, beef and non-lactating dairy cattle to provide 10 mg./lb. bodyweight daily.

BODYWEIGHT OF CATTLE	LEVEL OF CTC Per Head Per Day	FEEDING RATE Per Head Per Day
100 lb.	1 g	0.1 lb.
500 lb.	5 g	0.5 lb.

LIMITATIONS FOR USE: Feed for not more than 5 days at 10 mg/lb. per day rate. Feed approximately 400 g/ton varying with bodyweight and feed consumption to provide 10 mg/lb. per day. Calves under 250 lbs. fed at 10 mg/lb. per day rate must use in milk replacers or starter feeds.