



## DECOXY MEDICATED PELLET

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in ruminating and non-ruminating calves, including veal calves and cattle.

NET WEIGHT 50 LBS. (22.7kg)

### ACTIVE DRUG INGREDIENT

Decoquinatate ..... 0.125%  
(Equivalent to 568 mg/lb.)

### INGREDIENTS

Wheat Middlings, Soybean Hulls, Calcium Carbonate, Distillers Dried Grains/Solubles, Cane Molasses, Choice White Grease.

### FEEDING DIRECTIONS

Feed at a rate to provide 22.7 mg Decoquinatate (.04 lb. or .64 oz.) per 100 lbs. of bodyweight daily. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage. Do not use in feeds containing bentonite.

FEEDING RATE OF DECOXY .125%		
BWT.	LB./H/D	OZ./H/D
100 lbs.	.04 lbs.	0.64 oz.
250 lbs.	.10 lbs.	1.6 oz.
500 lbs.	.20 lbs.	3.2 oz.
1,000 lbs.	.40 lbs.	6.4 oz.

---

**WARNING:** Do not feed to cows producing milk for food.

---