



SUP'R PIG 15-25 DC MEDICATED

For treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella chloeraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline and for control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin.

NET WEIGHT 50 LBS.

ACTIVE DRUG INGREDIENTS

Chlortetracycline 400 grams/ton
 Tiamulin hydrogen fumerate 35 grams/ton

GUARANTEED ANALYSIS

Crude Protein	20.5% Min.
Lysine	1.35% Min.
Crude Fat	6.0% Min.
Crude Fiber	3.2% Max.
Calcium (Ca)	0.6% Min. ... 1.1% Max.
Phosphorus (P)	0.75% Min.
Selenium (Se)	0.3 ppm Min.
Zinc (Zn)	2,600 ppm Min.
Vitamin A	6,600 IU/lb. Min.

INGREDIENTS

Ground Yellow Corn, Soybean Meal, Dried Whey, Rolled Oat Groats, Fish Meal, Choice White Grease, Lactose, Dicalcium Phosphate, Calcium Carbonate, Hydrated Sodium Calcium Aluminosilicate, Zinc Oxide, L-Lysine, Sodium Selenite, DL-Methionine, Artificial Flavor, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Ethylenediamine Dihydriodide, Sodium Saccharin, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

DIRECTIONS FOR USE

Designed to be fed continuously as the sole ration for 14 days to pigs from 15 to 25 pounds bodyweight. Consult your PEET'S Representative for specific feeding recommendations.

CAUTION: Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g. Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds. Protect medicated feed from excessive temperatures. Do not store at warm temperatures, 104° F (40° C), for more than 7 days. Withdraw 2 days prior to slaughter.

LIMITATIONS FOR USE: Feed continuously as sole ration for 14 days. Not for use in swine weighing more than 250 pounds. Feed approximately 400 g/ton Chlortetracycline, varying with bodyweight and feed consumption, to provide 10 mg/lb. of bodyweight per day.