



CATTLE DEVELOPER RU1200 GRANULAR

A vitamin/mineral product formulated for growing calves and developing cattle to supplement rations containing high protein ingredients (soybean meal, ethanol by-products, haylage, early growth pasture and other protein sources). It works to improve feed efficiency for cattle being fed in confinement for slaughter.

NET WEIGHT 50 LBS. (22.7kg)

ACTIVE DRUG INGREDIENT

Monensin 1,200 g/ton

GUARANTEED ANALYSIS

Equivalent Crude Protein from Non-Protein Nitrogen	8.5% Min.
Calcium (Ca)	22.0% Min. .. 26.0% Max.
Phosphorus (P)	3.0% Min.
Salt (NaCl)	5.0% Min. .. 6.0% Max.
Magnesium (Mg)	1.3% Min.
Potassium (K)	2.5% Min.
Copper (Cu)	2,000 ppm Min.
Selenium (Se)	20 ppm Min.
Zinc (Zn)	4,000 ppm Min.
Vitamin A	300,000 IU/lb. Min.
Vitamin D-3	30,000 IU/lb. Min.
Vitamin E	300 IU/lb. Min.

INGREDIENTS

Calcium Carbonate, Dicalcium Phosphate, Distillers Dried Grains/Solubles, Salt, Monocalcium Phosphate, Potassium Chloride, Urea, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Sodium Selenite, Cane Molasses, Soybean Oil, Potassium/Magnesium Sulfate, Yeast Culture, Dried Aspergillus oryzae Fermentation Product, Ethylenediamine Dihydriodide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Choline Chloride, Niacin, Calcium Pantothenate, Riboflavin, Vitamin B-12 Supplement, Biotin, Menadione Dimethylpyrimidinol Bisulfite (Source of Vitamin K Activity), Thiamine, Pyridoxine Hydrochloride, Folic Acid, Ethoxyquin Preservative, Natural and Artificial Flavors, Artificial Color.

FEEDING DIRECTIONS

Feed 1/4 - 1/3 pound per head per day mixed in a grain mix or T.M.R. to provide 150 to 200 mg of Monensin and 50 mg thiamin daily. Do not free choice. Feed as the sole source of selenium. Consult your PEET'S Representative for ration balancing services.

WARNING: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating goats. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. This product should not be offered to sheep due to the copper level.

