



## FEEDLOT NATURAL DEC227 MEDICATED PELLET

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in ruminating and non-ruminating calves and veal calves.

NET WEIGHT 50 LBS. (22.7kg)

2 ton minimum

### ACTIVE DRUG INGREDIENT

Decoquinatate ..... 227 g/ton

### GUARANTEED ANALYSIS

Crude Protein	32.0% Min.
Crude Fat	2.0% Min.
Crude Fiber	10.0% Max.
Calcium (Ca)	7.0% Min. ... 8.0% Max.
Phosphorus (P)	0.85% Min.
Salt (NaCl)	2.5% Min. ... 3.5% Max.
Potassium (K)	1.5% Min.
Selenium (Se)	2.2 ppm Min.
Vitamin A	40,000 IU/lb. Min.
Vitamin D-3	4,000 IU/lb. Min.
Vitamin E	40 IU/lb. Min.

### INGREDIENTS

Soybean Meal, Calcium Carbonate, Distillers Dried Grains/Solubles, Pork Meat & Bone Meal, Cane Molasses, Salt, Linseed Meal, Dehydrated Alfalfa Meal, Wheat Middlings, Pellet Coating, Dicalcium Phosphate, Dried Aspergillus oryzae Fermentation Product, Potassium Chloride, Sodium Selenite, Manganese Sulfate, Zinc Sulfate, Ferrous Sulfate, Copper Sulfate, Cobalt Sulfate, Calcium Iodate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Ethoxyquin Preservative, Mineral Oil, Propionic Acid, Sorbic Acid, Acetic Acid, Benzoic Acid, Ammonium Hydroxide.

### FEEDING DIRECTIONS

Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Feed at the feeding rates listed below to provide 22.7 mg per 100 pounds of bodyweight of Decoquinatate daily for the prevention of coccidiosis in calves.

Calf Weight	LBS. to Feed Daily	DECOQUINATE provided (mg)
500	1.0	113.5
600	1.2	136.2
700	1.4	158.9
800	1.6	181.6
900	1.8	204.3

Consult your PEET'S Representative for ration balancing services to provide the optimum usage of grain and roughage and other medicated options available.

**WARNING:** Do not feed to cows producing milk for food. Do not use in feeds containing bentonite.

